





INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : G01N 33/566, 33/94, 33/60 C07D 471/14, 487/04		AI	(11) International Publication Number: WO 93/22681 (43) International Publication Date: 11 November 1993 (11.11.93)
(21) International Application Number: PCT/US93/03920 (22) International Filing Date: 30 April 1993 (30.04.93) (30) Priority data: 07/876,050 30 April 1992 (30.04.92) US (60) Parent Application or Grant (63) Related by Continuation US 07/876,050 (CON) Filed on 30 April 1992 (30.04.92) (71) Applicant (for all designated States except US): NEUROGEN CORPORATION [US/US]; 35 N.E. Industrial Road, Branford, CT 06405 (US).		(72) Inventors; and (75) Inventors/Applicants (for US only): SHAW, Kenneth [US/US]; 17 Cedar Hills Road, Weston, CT 06883 (US). HUTCHISON, Alan [US/US]; 175 Bartlett Drive, Madison, CT 06443 (US). THURKAUF, Andrew [US/US]; 6 Foxbridge Village Road, Branford, CT 06405 (US). TALLMAN, John [US/US]; 65 Prospect Avenue, Guilford, CT 06437 (US). (74) Agent: SARUSSI, Steven, J.; Allegretti & Witcoff, Ltd., Ten South Wacker Drive, Chicago, IL 60606 (US). (81) Designated States: AT, AU, BB, BG, BR, CA, CH, CZ, DE, DK, ES, FI, GB, HU, JP, KP, KR, KZ, LK, LU, MG, MN, MW, NL, NO, NZ, PL, RO, RU, SD, SE, SK, UA, US, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	
(54) Title: NOVEL GABA _A RECEPTOR SUBTYPES AND METHODS FOR SCREENING DRUG COMPOUNDS USING IMIDAZOQUINOXALINES AND PYRROLOPYRIMIDINES TO BIND TO GABA _A RECEPTOR SUBTYPES			
<div style="display: flex; justify-content: space-around; align-items: center;"> <div style="text-align: center;">  <p>(I)</p> </div> <div style="text-align: center;">  <p>(II)</p> </div> </div>			
(57) Abstract <p>The present invention provides methods for screening drug compounds utilizing compounds of formulas (I and II), and the pharmaceutically acceptable salts thereof where: R₁, R₂, R₃, and R₄ represent hydrogen, halogen, alkyl or alkoxy substituents; R₅ is hydrogen or lower alkyl; X and Y represent hydrogen, halogen, alkyl or alkoxy substituents; and Z is hydrogen or fluorine. The invention also provides tritium or iodine isotope radiolabeled compounds of the formulas (I and II) radiolabeled with tritium or isotopes of iodine. The invention further provides novel GABA_A receptor subtypes which specifically bind to compounds of formulas (I or II). The invention also provides GABA_A receptor subtypes which are bound <i>in situ</i> to a compound of formula (I or II). The compounds provided herein bind selectively to a novel subtype of the GABA_A binding site. Selective interaction of ligands at this unique receptor population results in pharmacological specificity which may lead to superior anxiolytics, cognition enhancers, anticonvulsants and sedative hypnotics.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NL	Netherlands
BE	Belgium	GN	Guinea	NO	Norway
BF	Burkina Faso	GR	Greece	NZ	New Zealand
BG	Bulgaria	HU	Hungary	PL	Poland
BJ	Benin	IE	Ireland	PT	Portugal
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SK	Slovak Republic
CI	Côte d'Ivoire	LJ	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	MC	Monaco	TG	Togo
DE	Germany	MG	Madagascar	UA	Ukraine
DK	Denmark	ML	Mali	US	United States of America
ES	Spain	MN	Mongolia	VN	Viet Nam
FI	Finland				

1.4-Benzodiazepines continue to be among the most widely used drugs in
35 the world. Principal among the benzodiazepines marketed are
chlordiazepoxide, diazepam, flurazepam, and triazolam. These compounds are
widely used as anxiolytics, sedative-hypnotics, muscle relaxants, and
anticonvulsants. A number of these compounds are extremely potent drugs;
such potency indicates a site of action with a high affinity and specificity for

individual receptors. Early electrophysiological studies indicated that a major action of benzodiazepines was enhancement of GABAergic inhibition. The benzodiazepines were capable of enhancing presynaptic inhibition of a monosynaptic ventral root reflex, a GABA-mediated event. (Schmidt et al., 1967, Arch. Exp. Path. Pharmacol. 258: 69-82). All subsequent electrophysiological studies (reviewed in Tallman et al. 1980, Science 207:274-81, Haefley et al., 1981, Handb. Exptl. Pharmacol. 33: 95-102) have generally confirmed this finding, and by the mid-1970s, there was a general consensus among electrophysiologists that the benzodiazepines could enhance the actions of GABA.

With the discovery of the "receptor" for the benzodiazepines and the subsequent definition of the nature of the interaction between GABA and the benzodiazepines, it appears that the behaviorally important interactions of the benzodiazepines with different neurotransmitter systems are due in a large part to the enhanced ability of GABA itself to modify these systems. Each modified system, in turn, may be associated with the expression of a behavior.

Studies on the mechanistic nature of these interactions depended on the demonstration of a high-affinity benzodiazepine binding site (receptor). Such a receptor is present in the CNS of all vertebrates phylogenetically newer than the boney fishes (Squires & Braestrup 1977, Nature 166: 732-34, Mohler & Okada, 1977, Science 198: 854-51, Mohler & Okada, 1977, Br. J. Psychiatry 133: 261-68). By using tritiated diazepam, and a variety of other compounds, it has been demonstrated that these benzodiazepine binding sites fulfill many of the criteria of pharmacological receptors; binding to these sites *in vitro* is rapid, reversible, stereospecific, and saturable. More importantly, highly significant correlations have been shown between the ability of benzodiazepines to displace diazepam from its binding site and activity in a number of animal behavioral tests predictive of benzodiazepine potency (Braestrup & Squires 1978, Br. J. Psychiatry 133: 249-60, Mohler & Okada, 1977, Science 198: 854-51, Mohler & Okada, 1977, Br. J. Psychiatry 133: 261-68). The average therapeutic doses of these drugs in man also correlate with potency at the GABA_A family of receptors (Tallman et al. 1980, Science 207: 274-281.). Thus, a radioreceptor assay, using tritiated diazepam, can serve as a tool for the discovery of novel therapeutics at the GABA_A receptor.

In 1978, it became clear that GABA and related analogs could interact at the low affinity (1 μ M) GABA binding site to enhance the binding of benzodiazepines to the clonazepam-sensitive site (Tallman et al. 1978, Nature, 274: 383-85). This enhancement was caused by an increase in the affinity of the benzodiazepine binding site due to occupancy of the GABA site. The data

were interpreted to mean that both GABA and benzodiazepine sites were allosterically linked in the membrane as part of a complex of proteins. For a number of GABA analogs, the ability to enhance diazepam binding by 50% of maximum and the ability to inhibit the binding of GABA to brain membranes
5 by 50% could be directly correlated. Enhancement of benzodiazepine binding by GABA agonists is blocked by the GABA receptor antagonist (+) bicuculline; the stereoisomer (-) bicuculline is much less active (Tallman et al., 1978, *Nature*, 274: 383-85).

Soon after the discovery of high affinity binding sites for the
10 benzodiazepines, it was discovered that a triazolopyridazine could interact with benzodiazepine receptors in a number of regions of the brain in a manner consistent with receptor heterogeneity or negative cooperativity. In these studies, Hill coefficients significantly less than one were observed in a number of brain regions, including cortex, hippocampus, and striatum. In cerebellum,
15 triazolopyridazine interacted with benzodiazepine sites with a Hill coefficient of 1 (Squires et al., 1979, *Pharma. Biochem. Behav.* 10: 825-30, Klepner et al. 1979, *Pharmacol. Biochem. Behav.* 11: 457-62). Thus, multiple benzodiazepine receptors were predicted in the cortex, hippocampus, striatum, but not in the cerebellum.

20 Based on these studies, extensive receptor autoradiographic localization studies were carried out at a light microscopic level. Although receptor heterogeneity has been demonstrated (Young & Kuhar 1980, *J. Pharmacol. Exp. Ther.* 212: 337-46, Young et al. 1981, *J. Pharmacol. Exp. Ther.* 216: 425-430, Niehoff et al. 1982, *J. Pharmacol. Exp. Ther.* 221: 670-75), no simple correlation between
25 localization of receptor subtypes and the behaviors associated with the region has emerged from the early studies. In addition, in the cerebellum, where one receptor was predicted from binding studies, autoradiography revealed heterogeneity of receptors (Niehoff et al., 1982, *J. Pharmacol. Exp. Ther.* 221: 670-75).

30 A physical basis for the differences in drug specificity for the two apparent subtypes of benzodiazepine sites was demonstrated by Sieghart & Karobath, 1980, *Nature* 286: 285-87. Using gel electrophoresis in the presence of sodium dodecyl sulfate, the presence of several molecular weight receptors for the benzodiazepines has been reported. The receptors were identified by
35 the covalent incorporation of radioactive flunitrazepam, a benzodiazepine which can covalently label all receptor types. The major labeled bands have molecular weights of 50,000 to 53,000, 55,000, and 57,000 and the triazolopyridazines inhibit labeling of the slightly higher molecular weight forms (53,000, 55,000, 57,000) (Sieghart et al. 1983, *Eur. J. Pharmacol.* 88: 291-99).

At that time, the possibility was raised that the multiple forms of the receptor represent "isoreceptors" or multiple allelic forms of the receptor (Tallman & Gallager 1985, *Ann. Rev. Neurosci.* 8, 21-44). Although common for enzymes, genetically distinct forms of receptors have not generally been described. As we begin to study receptors using specific radioactive probes and electrophoretic techniques, it is almost certain that isoreceptors will emerge as important in investigations of the etiology of psychiatric disorders in people.

The GABA_A receptor subunits have been cloned from bovine and human cDNA libraries (Schofield et al. 1988, *Nature* 328, 221-227; Garrett et al., *BBRC* 156, 1039-1045, 1989). A number of distinct cDNAs were identified as subunits of the GABA_A receptor complex by cloning and expression. These are categorized into α , β , γ , δ , ϵ , and provide a molecular basis for the GABA_A receptor heterogeneity and distinctive regional pharmacology (Shivers et al. 1990, *Neuron* 4, 919-928; Levitan et al. 1988, *Nature* 335, 76-79). The γ subunit appears to enable drugs like benzodiazepines to modify the GABA responses (Pritchett et al. 1989, *Nature* 338, 582-585). The delta subunit is associated with GABA_A receptors that do not respond to benzodiazepines but may respond to other related compounds (Shivers et al., 1989, op. cit.) The presence of low Hill coefficients in the binding of ligands to the GABA_A receptor indicates unique profiles of subtype specific pharmacological action.

Drugs that interact at the GABA_A receptor can possess a spectrum of pharmacological activities depending on their abilities to modify the actions of GABA. For example, the beta-carbolines were first isolated based upon their ability to inhibit competitively the binding of diazepam to its binding site (Nielsen et al., 1979, *Life Sci.* 25: 679-86). Although predictive about potency at a receptor, the receptor binding assay does not totally predict the precise biological activity of such compounds; agonists, partial agonists, inverse agonists, and antagonists can inhibit binding. When the beta-carboline structure was determined, it was possible to synthesize a number of analogs and test these compounds behaviorally. It was immediately realized that the beta-carbolines could antagonize the actions of diazepam behaviorally (Tenen & Hirsch, 1980, *Nature* 288: 609-10). In addition to this antagonism, beta-carbolines possess intrinsic activity of their own opposite to that of the benzodiazepines; they become known as inverse agonists.

In addition, a number of other specific antagonists of the benzodiazepine receptor were developed based on their ability to inhibit the binding of benzodiazepines. The best studied of these compounds is an imidazodiazepine, (Hunkeler et al., 1981, *Nature* 290: 514-516). This compound is a high affinity competitive inhibitor of benzodiazepine and beta-carboline

binding and is capable of blocking the pharmacological actions of both these classes of compounds. By itself, it possesses little intrinsic pharmacological activity in animals and humans (Hunkeler et al., 1981, *Nature* 290: 514-16; Darragh et al., 1983, *Eur. J. Clin. Pharmacol.* 14: 569-70). When a radiolabeled
5 form of this compound was studied (Mohler & Richards, 1981, *Nature* 294: 763-65), it was demonstrated that this compound would interact with the same number of sites as the benzodiazepines and beta-carbolines, and that the interactions of these compounds were purely competitive. This compound was the ligand of choice for binding to GABA_A receptors to discover older leads
10 because it does not possess receptor subtype specificity and measures each state of the receptor.

The study of the interactions of a wide variety of compounds similar to the above has led to the categorizing of these compounds. Presently, those
15 compounds possessing activity similar to the benzodiazepines are called agonists. Compounds possessing activity opposite to benzodiazepines are called inverse agonists, and the compounds blocking both types of activity have been termed antagonists. This categorization has been developed to emphasize the fact that a wide variety of compounds can produce a spectrum of
20 pharmacological effects, to indicate that compounds can interact at the same receptor to produce opposite effects, and to indicate that beta-carbolines and antagonists with intrinsic anxiogenic effects are not synonymous. A biochemical test for the pharmacological and behavioral properties of compounds that interact with the benzodiazepine receptor continues to
25 emphasize the interaction with the GABAergic system. In contrast to the benzodiazepines, which show an increase in their affinity due to GABA (Tallman et al., 1978, *Nature* 274: 383-85, Tallman et al., 1980, *Science* 207: 274-81), compounds with antagonist properties show little GABA shift (i.e., change in receptor affinity due to GABA) (Mohler & Richards 1981, *Nature* 294: 763-65),
30 and the inverse agonists actually show a decrease in affinity due to GABA [(Braestrup & Nielson 1981, *Nature* 294: 472-474)]. Thus, the GABA shift in a receptor binding assay in addition to high affinity predicts generally the expected behavioral properties of the compounds.

SUMMARY OF THE INVENTION

The present invention provides methods for screening drug compounds comprising the steps of:

5

- (a) conducting in a reaction mixture a competition reaction between compounds of formulas I or II radiolabeled with tritium or an iodine isotope and the drug to be tested, for a GABA_A receptor having a binding site for the compound of formula I or II;

10


- (c) separating the GABA_A receptor bound radiolabeled compound of formula I or II from the reaction mixture; and

15


- (b) measuring the radioactivity resulting from specific binding of the radiolabeled compound of formula I or II to the GABA_A receptor.

where formulas I and II are

20



I



II

and the pharmaceutically acceptable salts thereof
where:

25

R₁, R₂, R₃, and R₄ are the same or different and represent hydrogen, halogen, straight chain or branched lower alkyl having 1-6 carbon atoms, or straight chain or branched lower alkoxy having 1-6 carbon atoms;


30

R5 is hydrogen or straight chain or branched lower alkyl having 1-6 carbon atoms;


5 X and Y are the same or different and represent hydrogen, halogen, straight chain or branched lower alkyl having 1-6 carbon atoms, or straight chain or branched lower alkoxy having 1-6 carbon atoms; and

Z is hydrogen or fluorine.

10 The invention also provides tritium or iodine isotope radiolabeled compounds of the formulas



and



15

and the pharmaceutically acceptable salts thereof
where:

20 R1, R2, R3, R4 are the same or different and represent hydrogen, halogen, straight chain or branched lower alkyl having 1-6 carbon atoms, or straight chain or branched lower alkoxy having 1-6 carbon atoms;

25 R5 is hydrogen or straight chain or branched lower alkyl having 1-6 carbon atoms;

X and Y are the same or different and represent hydrogen, halogen, straight chain or branched lower alkyl having 1-6 carbon atoms, or straight chain or branched lower alkoxy having 1-6 carbon atoms; and

Z is hydrogen or fluorine.

Further, the invention provides novel GABA_A receptor subtypes which
5 specifically bind to compounds of formulas I or II.

The invention also provides GABA_A receptor subtypes which are bound
in situ to a compound of formula I or II.

The invention provides compounds and methods which can be utilized to
discover and develop novel therapeutic agents useful in treating anxiety, sleep
10 and seizure disorders and enhancing alertness. The compounds provided
herein bind selectively to a novel subtype of the GABA_A binding site. This
novel subtype is a subset of GABA_A receptors which is distinct from those
GABA_A receptors disclosed in the prior art. Selective interaction of ligands at
this unique receptor population results in pharmacological specificity which
15 may lead to superior anxiolytics, cognition enhancers, anticonvulsants and
sedative hypnotics.

BRIEF DESCRIPTION OF THE DRAWING



Figures 1A-G show representative imidazoquinoxalines and imidazopyrimidines which are employed in the methods of the present invention.

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides methods for screening drug compounds comprising the steps of:

- 5 (a) conducting in a reaction mixture a competition reaction between compounds of formulas I or II radiolabeled with tritium or an iodine isotope and the drug to be tested, for a GABA_A receptor having a binding site for the compound of formula I or II;
- 10 (c) separating the GABA_A receptor bound radiolabeled compound of formula I or II from the reaction mixture; and
- 15 (b) measuring the radioactivity resulting from specific binding of the radiolabeled compound of formula I or II to the GABA_A receptor,

where formulas I and II are



and the pharmaceutically acceptable salts thereof
where:

- 25 R₁, R₂, R₃, and R₄ are the same or different and represent hydrogen, halogen, straight chain or branched lower alkyl having 1-6 carbon atoms, or straight chain or branched lower alkoxy having 1-6 carbon atoms;


-11-

R5 is hydrogen or straight chain or branched lower alkyl having 1-6 carbon atoms;


5 X and Y are the same or different and represent hydrogen, halogen, straight chain or branched lower alkyl having 1-6 carbon atoms, or straight chain or branched lower alkoxy having 1-6 carbon atoms; and

Z is hydrogen or fluorine.

10 The invention also provides tritium or iodine isotope radiolabeled compounds of the formulas



and



15

and the pharmaceutically acceptable salts thereof
where:

20 R1, R2, R3, R4 are the same or different and represent hydrogen, halogen, straight chain or branched lower alkyl having 1-6 carbon atoms, or straight chain or branched lower alkoxy having 1-6 carbon atoms;

25 R5 is hydrogen or straight chain or branched lower alkyl having 1-6 carbon atoms;

X and Y are the same or different and represent hydrogen, halogen, straight chain or branched lower alkyl having 1-6 carbon atoms, or straight chain or branched lower alkoxy having 1-6 carbon atoms; and

Z is hydrogen or fluorine.

Further, the invention provides novel GABA_A receptor subtypes which
5 specifically bind to compounds of formulas I or II.

The invention also provides GABA_A receptor subtypes which are bound
in situ to a compound of formula I or II.

Representative compounds suitable for use in the drug screening
methods of the present invention, which are encompassed by formulas I and II
10 include, but are not limited to, the compounds in Figures 1A-G and their
pharmaceutically acceptable salts.

The compounds which may be employed in the inventive methods may
be radiolabeled with any convenient radiolabel. Such radiolabels are, for
example, isotopes of hydrogen, carbon, and iodine. A convenient hydrogen
15 isotope is tritium. The compounds may also be prepared to include various
isotopes of iodine, such as, for example, ¹²⁷I. The suitable isotopes of carbon
are ¹³C and ¹⁴C.

The invention provides compounds and methods which can be utilized to
discover and develop novel therapeutic agents useful in treating anxiety, sleep
20 and seizure disorders and enhancing alertness. The compounds provided
herein bind selectively to a novel subtype of the GABA_A binding site. This
novel subtype is a subset of GABA_A receptors which is distinct from those
GABA_A receptors disclosed in the prior art. Selective interaction of ligands at
this unique receptor population results in pharmacological specificity which
25 may lead to superior anxiolytics, cognition enhancers, anticonvulsants and
sedative hypnotics.


Prior art compounds known to have high affinity for all GABA_A
receptors did not exhibit such affinity for the receptor of the present
invention. The prior art compounds all demonstrated low affinity towards the
30 novel receptor subtype of the invention. Table II below shows the affinities of
prior art compounds towards known GABA_A receptors. These data indicate that
one skilled in the art would have expected those compounds to have similar
affinities towards the GABA_A receptor of the present invention. The data
presented in Tables I and II clearly indicate that the prior art compounds have
35 dramatically lower affinities towards the novel receptor subtype.

An illustration of the preparation of compounds of the present
invention is given in Schemes I and II. Those having skill in the art will
recognize that the starting materials may be varied and additional steps

employed to produce either tritiated or iodinated compounds encompassed by the present invention, as demonstrated by the following examples.


Scheme I

5



-14-

Scheme II



5

wherein:

-15-

R₁, R₂, R₃, and R₄ are the same or different and represent hydrogen, halogen, straight chain or branched lower alkyl having 1-6 carbon atoms, or straight chain or branched lower alkoxy having 1-6 carbon atoms;

5

R₅ is hydrogen or straight chain or branched lower alkyl having 1-6 carbon atoms;

10

X and Y are the same or different and represent hydrogen, halogen, straight chain or branched lower alkyl having 1-6 carbon atoms, or straight chain or branched lower alkoxy having 1-6 carbon atoms; and

15

Z is hydrogen or fluorine.

The methods for drug screening of the present invention are competitive assays where two compounds, one of which is radiolabeled, compete for a GABA_A receptor. In these methods, either of the compounds of formulas I and II may be radiolabeled or the drug to be tested may be radiolabeled. Preferred methods are those in which a compound of formula I or II is radiolabeled and allowed to compete for a GABA_A receptor.

The invention is illustrated further by the following examples which are not to be construed as limiting the invention in scope or spirit to the specific procedures and compounds described in them.

EXAMPLE 1

Drug Screening Assay

The methods for screening drug compounds of the invention using the
5 novel GABA_A receptor are illustrated by the following assays for GABA_A
receptor subtype activity.

- Assays were carried out using the tissue preparation described in
Thomas and Tallman (J. Bio. Chem. 156: 9838-9842, J. Neurosci. 3:433-440, 1983).
- 10 Rat cortical tissue was dissected and homogenized in 25 volumes (w/v) of
0.05 M Tris HCl buffer (pH 7.4 at 4°C). The tissue homogenate was centrifuged
in the cold (4°C) at 20,000 x g for 20'. The supernatant was decanted and the
pellet was rehomogenized in the same volume of buffer and again centrifuged
at 20,000 x g. The supernatant was decanted and the pellet was frozen at -20°C
15 overnight. The pellet was then thawed and rehomogenized in 25 volume
(original wt/vol) of buffer and the procedure was carried out twice. The pellet
was finally resuspended in 50 volumes (w/vol) of 0.05 M Tris HCl buffer
containing 1uM Ro 15-1788, 1uM sodium ascorbate and 10uM muscimol (pH 7.4
at 4°C).
- 20 Incubation mixtures contained 400 µl of tissue homogenate, 5 µl of
100nM ligand in above buffer [final ligand concentration was 0.5 nM ³H-
Compound 1 (specific activity 67 Ci/mmol)], drug¹ or blocker and buffer to a
total volume of 1 mL. Incubations were carried for 20 min at 4°C then were
rapidly filtered through GFB filters to separate free and bound ligand. Filters
25 were washed twice with fresh 0.05 M Tris HCl buffer (pH 7.4 at 4°C) and
counted in a liquid scintillation counter. 10 µM Compound 3 was added to some
tubes to determine nonspecific binding. Data were collected in triplicate
determinations, averaged and % inhibition of total specific binding was
calculated. Total Specific Binding = Total - Nonspecific. Typically the specific
30 binding represented about 75 to 85% of total binding. In some cases, the
amounts of unlabeled drugs was varied and total displacement curves of
binding were carried out. Data were converted to a form suitable for the
calculation of IC₅₀ and Hill Coefficient (n_H). These IC 50's were converted to
Ki's using the Cheng Prusoff equation (Cheng and Prusoff, 1973, Biochem.
35 Pharmacol. 22, 3099.). Data for the compounds of this invention as well as
certain reference compounds are listed in Table I. All of the reference prior
art compounds tested demonstrated low affinity for this receptor subtype.

TABLE I

	<u>Compound Name or Number</u> ¹	<u>Ki (nM)</u>
5	Alprazolam	8000
	Diazepam	5000
	Flumazenil	7500
	Bretazenil	15000
10	Zolpidem	2200
	Compound 1	0.5
	Compound 2	3.9
	Compound 3	0.9
	Compound 4	0.8
15	Compound 5	6
	Compound 6	0.55
	Compound 7	50

1 Compound numbers relate to compounds of the present invention
 20 shown in Figures 1A-G.


Table II lists the affinities of prior art compounds for known GABA_A receptors.

Table II

	<u>Compound Name</u>	<u>Ki (nM)</u>
25	Diazepam	6
	Flumazenil	1
30	Bretazenil	1
	Zolpidem	20

Examples II-XIII are syntheses of the compounds of the invention.

EXAMPLE II




5


To a solution of 2-Nitrophenyl isocyanate (3.34 g) in 100 mL of toluene was added 3-Bromo-4-ethoxyaniline (3.1 g). The mixture was stirred at 20°C for 30 min. Hexane (300 mL) was added and the resulting solid was filtered and
10 dried to yield N-(2-Nitrophenyl)-N'-(3-bromo-4-ethoxyphenyl)-urea as a light yellow solid.

EXAMPLE III

15




A solution containing N-(2-Nitrophenyl)-N'-(3-bromo-4-ethoxyphenyl)-urea (5.76 g) and chloroacetyl chloride (40 mL) was refluxed under nitrogen
20 for 30 min. After the excess chloroacetyl chloride was removed *in vacuo*, diethyl ether (50 mL) was added and the resulting solid was filtered and dried to yield N'-(2-chloroacetyl)-N-(2-nitrophenyl)-N'-(3-bromo-4-ethoxyphenyl)-urea as a white solid.


EXAMPLE IV

5 A solution of N'-(2-Chloroacetyl)-N-(2-nitrophenyl)-N'-(3-bromo-4-ethoxyphenyl)-urea (3.7 g), dimethylformamide (15 mL) and diisopropyl-ethylamine (15 mL) was refluxed for 5 min. The hot mixture was allowed to cool to room temperature and precipitated by adding the mixture to 200 mL of water. The precipitate was collected and dried to yield 1-(2-Nitrophenyl)-3-(3-bromo-4-ethoxyphenyl)-imidazolidine-2,4(1H,3H)-dione.


10

EXAMPLE V

5 To a solution containing 1-(2-Nitrophenyl)-3-(3-bromo-4-ethoxy-phenyl)-imidazoline-2,4(1H,3H)-dione (1.18 g) in anhydrous methylene chloride (5 mL) under nitrogen was added tris(dimethylamino)methane (1 mL). The reaction was stirred at room temperature for 20 min and the solvent was removed *in vacuo*. To the resulting oil was added iron powder (5 g) and acetic
10 acid (250 mL). This mixture was carefully heated to reflux for 3 min followed by stirring the reaction for an additional 30 min. The heterogeneous mixture was diluted with 10% methanol-methylene chloride (200 mL) and filtered through silica gel using 10% methanol/methylene chloride as eluant. The solvent was removed *in vacuo* and hot ethanol (200 mL) was added. To this mixture was
15 added water (200 mL) and the resulting solid was filtered and washed successively with ethanol, ethyl acetate, diethyl ether and dried to yield 2-(3-Bromo-4-ethoxyphenyl)-imidazo[1,5-a]quinoxaline-1,3(2H,5H)-dione as a yellow solid.


EXAMPLE VI

5 To 2-(3-Bromo-4-ethoxyphenyl)-imidazo[1.5.a]quinoxaline-1.3(2H.5H)-
dione (250 mg) in dioxane (10 mL) was added Bromine (1 mL). The reaction
mixture was heated to 80°C for 1.5 hr and then poured directly into a
suspension of zinc dust (2.0 gm) in acetic acid (50 mL). The reaction mixture
was heated to reflux for 5 min and then allowed to cool to room temperature
10 over 45 min. The suspension was filtered through celite and evaporated under
reduced pressure. The resulting solid was stirred for 10 min with a solution
containing ethanol (25 mL) and water (50 mL) and filtered. The solid was
washed with ethanol (10 mL) followed by ether (10 mL) to yield 6,8-Dibromo-2-
(3-Bromo-4-ethoxyphenyl)-imidazo[1.5.a]quinoxaline-1.3(2H.5H)-dione as a
15 yellow solid.

EXAMPLE VII


Compound 1

5
10
15
6,8-Dibromo-2-(3-Bromo-4-ethoxyphenyl)-imidazo[1,5,a]quinoxaline-1,3(2H,5H)-dione (7.5 mg) dissolved in a suspension containing dimethylformamide (1 mL), absolute ethanol (1mL), triethylamine (10 ul) and 10% palladium on carbon (5 mg) was stirred under tritium gas at atmospheric pressure for 30 min. The reaction vessel was degassed, filtered through celite and the solvent was removed at reduced pressure. The reaction mixture was then subjected to three ethanol (3 mL) and one acetic acid (3 mL) dissolution-evaporation cycles to eliminate all radioactive exchangeable protons. The desired tritiated 2-(4-Ethoxyphenyl)imidazo[1,5,a]quinoxaline-1,3(2H,5H)-dione (Compound 1) was purified by preparative HPLC. It had Specific Activity of 67 Ci/mmol.


EXAMPLE VIII

- 5 A mixture of 2-Fluoro-5-methoxybenzimidine (10.3 g) and dimethyl malonate (8.09 g) in dry dimethyl sulfoxide (7 mL) was allowed to stand at room temperature for 24 h.. The precipitated product was collected and washed with water and ether to afford 2-Phenyl-4,6-dihydroxy-pyrimidine as a white solid.


10

EXAMPLE IX


- 15 To a suspension of 2-(2-Fluoro-5-methoxyphenyl)-4,6-dihydroxy-pyrimidine (12 g) in 35 mL of acetic acid is added 12 mL of 90% nitric acid and the mixture is heated at 50°C for 45 min. The reaction mixture is diluted with 150 mL of water and the product is collected, washed with water and ethanol and oven dried to afford 2-(2-Fluoro-5-methoxyphenyl)-5-nitro-4,6-dihydroxy-
20 pyrimidine as a pink solid.

Example X

- 5 A mixture of 2-(2-Fluoro-5-methoxyphenyl)-5-nitro-4,6-dihydroxy-pyrimidine (10 g), diethylaniline (7 g) and phosphorus oxychloride (100 mL) was heated at reflux for 40 min. The reaction mixture was concentrated *in vacuo* and the residue was partitioned between 50% ether in ethyl acetate and water. The organic layer was dried over magnesium sulfate and the solvent was
- 10 removed *in vacuo*. The residue was filtered through silica gel with ether/methylene chloride as the eluent to afford 2-(2-Fluoro-5-methoxyphenyl)-5-nitro-4,6-dichloro-pyrimidine as a yellow solid.

Example XI

- 5 A mixture of cyclohexanone (98 mg) and pyrrolidine (71 mg) and 4A molecular sieves (500 mg) in 1 mL of benzene is allowed to stand at room temperature until enamine formation was complete (ca. 16 h). The resulting solution of enamine was cannulated into a solution of 2-(2-Fluoro-5-methoxyphenyl)-5-nitro-4,6-dichloro-pyrimidine (300 mg) and
- 10 diisopropylethylamine (129 mg) in 5 mL of methylene chloride. After 30 min at room temperature the reaction mixture was concentrated *in vacuo* and treated with 3 mL of 3N HCl and 3 mL of ethanol. The reaction mixture was concentrated again and the residue was subjected to flash chromatography on silica gel with 20% ethyl acetate/hexane as the eluent to afford 2-[4-(2-(2-
- 15 Fluoro-5-methoxyphenyl)-5-nitro-6-chloro-pyrimidinyl)]-cyclohexan-1-one as a white solid.

EXAMPLE XII

(Compound 2)

5

A mixture of 2-[4-(2-(2-Fluoro-5-methoxyphenyl)-5-nitro-6-chloro-pyrimidinyl)]-cyclohexan-1-one (280 mg), triethylamine (300 mg) and 10% Pd/C catalyst (25 mg) in 10 mL of ethanol was hydrogenated under 1 atmosphere of hydrogen at room temperature for 16 h. After filtration
10 through celite the solvent was removed *in vacuo* and the residue was subjected to flash chromatography on silica gel with 50% ethyl acetate /hexane as the eluent to afford 2-(2-Fluoro-5-methoxyphenyl)-6,7,8,9-tetrahydro-5H-indolo[3,2-d]-pyrimidine melting at 197-198°C (Compound 2) after trituration with hexane/ether. Performing essentially the same procedure with tritium
15 gas afforded tritiated 2-(2-Fluoro-5-methoxyphenyl)-6,7,8,9-tetrahydro-5H-indolo[3,2-d]-pyrimidine (Compound 2).

EXAMPLE XIII

20

The following compounds were prepared essentially according to the procedures described in Examples I-XI.

- a) 2-(4-Propyloxyphenyl)imidazo[1,5,a]quinoxaline-1,3(2H,5H)-dione
25 (Compound 3).
- b) 2-(4-Ethoxyphenyl)-7-methyl-imidazo[1,5,a]quinoxaline-1,3(2H,5H)-dione (Compound 4).
- 30 c) 2-(4-Ethoxyphenyl)-5-methyl-imidazo[1,5,a]quinoxaline-1,3(2H,5H)-dione (Compound 5).



- d) 2-(3-Methoxyphenyl)imidazo[1,5.a]quinoxaline-1,3(2H,5H)-dione
(Compound 6).
- e) 2-(3-Methoxyphenyl)-6,7,8,9-tetrahydro-5H-indolo[3,2-d]-pyrimidine
5 (Compound 7).

The invention and the manner and process of making and using it, are now described in such full, clear, concise and exact terms as to enable any person skilled in the art to which it pertains, to make and use the same. It is to
10 be understood that the foregoing describes preferred embodiments of the present invention and that modifications may be made therein without departing from the spirit or scope of the present invention as set forth in the claims. To particularly point out and distinctly claim the subject matter regarded as invention, the following claims conclude this specification.

WHAT IS CLAIMED IS:

1. A method for drug screening comprising the steps of:
 - 5 (a) conducting in a reaction mixture a competition reaction between compounds of formulas I or II radiolabeled with tritium or an iodine isotope and the drug to be tested, for a GABA_A receptor having a binding site for the compound of formula I or II;
 - 10 (c) separating the GABA_A receptor bound radiolabeled compound of formula I or II from the reaction mixture; and
 - (b) measuring the radioactivity resulting from specific binding of the radiolabeled compound of formula I or II to the GABA_A receptor,

where formulas I and II are



and the pharmaceutically acceptable salts thereof
where:


- 25 R₁, R₂, R₃, R₄ are the same or different and represent hydrogen, halogen, straight chain or branched lower alkyl having 1-6 carbon atoms, or straight chain or branched lower alkoxy having 1-6 carbon atoms;

R5 is hydrogen or straight chain or branched lower alkyl having 1-6 carbon atoms;


5 X and Y are the same or different and represent hydrogen, halogen, straight chain or branched lower alkyl having 1-6 carbon atoms, or straight chain or branched lower alkoxy having 1-6 carbon atoms; and

Z is hydrogen or fluorine.

10 2. Tritium or iodine isotope radiolabeled compounds of the formulas



and



15 and the pharmaceutically acceptable salts thereof
where:

20 R1, R2, R3, R4 are the same or different and represent hydrogen, halogen, straight chain or branched lower alkyl having 1-6 carbon atoms, or straight chain or branched lower alkoxy having 1-6 carbon atoms;



R5 is hydrogen or straight chain or branched lower alkyl having 1-6 carbon atoms;

25

X and Y are the same or different and represent hydrogen, halogen, straight chain or branched lower alkyl having 1-6 carbon atoms, or straight chain or branched lower alkoxy having 1-6 carbon atoms; and

Z is hydrogen or fluorine.

3. A GABA_A receptor which specifically binds to compounds of
5 formulas I or II, where formulas I and II are:



10 and the pharmaceutically acceptable salts thereof
where:

15 R₁, R₂, R₃, R₄ are the same or different and represent hydrogen, halogen, straight chain or branched lower alkyl having 1-6 carbon atoms, or straight chain or branched lower alkoxy having 1-6 carbon atoms;

R₅ is hydrogen or straight chain or branched lower alkyl having 1-6 carbon atoms;

20 X and Y are the same or different and represent hydrogen, halogen, straight chain or branched lower alkyl having 1-6 carbon atoms, or straight chain or branched lower alkoxy having 1-6 carbon atoms; and

25 Z is hydrogen or fluorine.

4. The GABA_A receptor, of Claim 4, which is bound *in situ* to a compound of formula I or II.

5. A method according to Claim 1, where the compound is 2-(4-Ethoxyphenyl)imidazo[1,5,a]quinoxaline-1,3(2H,5H)-dione.
6. A method according to Claim 1, where the compound is 2-(2-Fluoro-5-methoxyphenyl)-6,7,8,9-tetrahydro-5H-indolo[3,2-d]-pyrimidine.
7. A method according to Claim 1, where the compound is 2-(4-Propyloxyphenyl)imidazo[1,5,a]quinoxaline-1,3(2H,5H)-dione.
8. A method according to Claim 1, where the compound is 2-(4-Ethoxyphenyl)-7-methyl-imidazo[1,5,a]quinoxaline-1,3(2H,5H)-dione.
9. A method according to Claim 1, where the compound is 2-(4-Ethoxyphenyl)-5-methyl-imidazo[1,5,a]quinoxaline-1,3(2H,5H)-dione.
10. A method according to Claim 1, where the compound is 2-(3-Methoxyphenyl)imidazo[1,5,a]quinoxaline-1,3(2H,5H)-dione.
11. A method according to Claim 1, where the compound is 2-(3-Methoxyphenyl)-6,7,8,9-tetrahydro-5H-indolo[3,2-d]-pyrimidine.
12. A GABAA receptor according to Claim 3 which specifically binds to the compound 2-(4-Ethoxyphenyl)imidazo[1,5,a]quinoxaline-1,3(2H,5H)-dione.
13. A GABAA receptor according to Claim 3 which specifically binds to the compound 2-(2-Fluoro-5-methoxyphenyl)-6,7,8,9-tetrahydro-5H-indolo[3,2-d]-pyrimidine.
14. A GABAA receptor according to Claim 3 which specifically binds to the compound 2-(4-Propyloxyphenyl)imidazo[1,5,a]quinoxaline-1,3(2H,5H)-dione.
15. A GABAA receptor according to Claim 3 which specifically binds to the compound 2-(4-Ethoxyphenyl)-7-methyl-imidazo[1,5,a]quinoxaline-1,3(2H,5H)-dione.
16. A GABAA receptor according to Claim 3 which specifically binds to the compound 2-(4-Ethoxyphenyl)-5-methyl-imidazo[1,5,a]quinoxaline-1,3(2H,5H)-dione.

17. A GABA_A receptor according to Claim 3 which specifically binds to the compound 2-(3-Methoxyphenyl)imidazo[1,5,a]quinoxaline-1,3(2H,5H)-dione.

5

18. A GABA_A receptor according to Claim 3 which specifically binds to the compound 2-(3-Methoxyphenyl)-6,7,8,9-tetrahydro-5H-indolo[3,2-d]-pyrimidine.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 93/03920

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all) ⁶		
According to International Patent Classification (IPC) or to both National Classification and IPC		
Int.Cl. 5 G01N33/566; G01N33/94; G01N33/60; C07D471/14 C07D487/04		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
Int.Cl. 5	G01N ; C07D ; C07B	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁸		
III. DOCUMENTS CONSIDERED TO BE RELEVANT⁹		
Category ¹⁰	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
A	WO,A,9 206 094 (NEUROGEN CORPORATION) 16 April 1992 see the whole document ---	1-4, 11, 13, 18
P,A	WO,A,9 207 853 (NEUROGEN CORPORATION) 14 May 1992 see the whole document ---	1-10, 12, 14-17
A	NATURE. vol. 328, 16 July 1987, LONDON GB pages 221 - 227 PETER R. SCHOFIELD ET AL. 'Sequence and functional expression of the GABA _A receptor shows a ligand gated receptor super-family' cited in the application see the whole document --- -/--	1-18
<p>¹⁰ Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"A" document member of the same patent family</p>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search 24 AUGUST 1993		Date of Mailing of this International Search Report 06.09.93
International Searching Authority EUROPEAN PATENT OFFICE		Signature of Authorized Officer DÖPFER K.P.

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No.
A	NATURE. vol. 294, 24 December 1981, LONDON GB pages 763 - 765 H. MÖHLER AND J.G. RICHARDS 'Agonist and antagonist benzodiazepine receptor interaction in vitro' cited in the application see the whole document ----	1-18
A	NATURE. vol. 335, 1 September 1988, LONDON GB pages 76 - 79 EDWIN S. LEVITAN 'Structural and functional basis for GABA _A receptor heterogeneity' cited in the application see the whole document -----	1-18
A	NATURE. vol. 286, 17 July 1980, LONDON GB pages 285 - 287 WERNER SIEGHART AND MANFRED KAROBATH 'Molecular heterogeneity of benzodiazepine receptors' cited in the application -----	1-18

**ANNEX TO THE INTERNATIONAL SEARCH REPORT
ON INTERNATIONAL PATENT APPLICATION NO.**

US 9303920
SA 74068

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

24/08/93

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9206094	16-04-92	AU-A- 8732691	28-04-92
		CA-A- 2091986	10-04-92
		EP-A- 0552237	28-07-93
		US-A- 5216159	01-06-93

WO-A-9207853	14-05-92	US-A- 5130430	14-07-92
		EP-A- 0555391	18-08-93

EPO FORM P0079